arXiv:1410.0137v1 [hep-ph] 1 Oct 2014

$\bar{D}^* D_0^*$ and $\bar{B}^* B_0^*$ (1^{--}) molecules at N2LO from QSSR*

F. Fanomezana[a,**], S. Narison[b], Andry Rabemananjara[a]

[a] *Institute of High-Energy Physics of Madagascar(iHEPMAD),University of Antananarivo, Madagascar*
[b] *Laboratoire Univers et Particules (LUPM), CNRS-IN2P3 & Université de Montpellier II, Case 070, Place Eugène Bataillon, 34095 - Montpellier Cedex 05, France.*

Abstract

We estimate the $\bar{D}^* D_0^*$ and $\bar{B}^* B_0^*$ (1^{--}) molecules masses and couplings using QCD spectral sum rules (QSSR) known perturbatively to N2LO of PT series and including the contributions of non-perturabtive condensates up to the dimension-eight. Our results improve earlier LO results obtained from QSSR in the current literature. We obtain $M_{barD^* D_0^*} = 5244(228)$ MeV which is heavier than the experimental candidates $Y(4260)$, $Y(4360)$, $Y(4660)$ suggesting that they cannot be pure molecule states. We predict $M_{\bar{B}^* B_0^*} = 11920(159)$ MeV to be tested in $B-$factory experiments.

Keywords: QCD spectral sum rules, molecules states, heavy quarkonia

1. Introduction

Motivated by the recently observed 1^{--} states Y(4260), Y(4360), Y(4660) from their decays to $J/\psi\pi\pi$ decays [1] often interpreted as molecules or four-quark states [2], we improve our previous results [3] obtained to lowest order (LO) of PT series from the QCD spectral sum rules [4, 5] by evaluating the mass and coupling of the $\bar{D}^* D_0^*$ and $\bar{B}^* B_0^*$ (1^{--}) molecules at N2LO of PT QCD and including up to dimension 8 condensates.

2. QCD analysis of spin one molecule

• *Current and two-point fonction*
The two-point correlation function associated to the $\bar{D}^* D_0^*$ and $\bar{B}^* B_0^*$ (1^{--}) molecule current is defined as:

$$\begin{aligned}\Pi^{\mu\nu}_{mol}(q) &= i\int d^4x\, e^{iq.x}\langle 0|T J^{\mu}(x) J^{\nu\dagger}(0)|0\rangle \\ &= -(q^2 g^{\mu\nu} - q^\mu q^\nu)\Pi_{mol}(q^2) \\ &\quad + q^\mu q^\nu \Pi^{(0)}_{mol}(q^2)\,, \end{aligned} \tag{1}$$

with the current:

$$\begin{aligned} J^\mu &\equiv (\bar{Q}\gamma^\mu q)(\bar{q}Q)\,, \\ Q &\equiv c, b \quad \text{and} \quad q \equiv u, d\,, \end{aligned} \tag{2}$$

where Π_{mol} and $\Pi^{(0)}_{mol}$ are respectively associated to the spin 1 and 0 molecule states. Due to its analyticity property, the correlation function $\Pi_{mol}(q^2)$, obeys the dispersion relation:

$$\Pi_{mol}(q^2) = \frac{1}{\Pi}\int_{4m_c^2}^{\infty} ds \frac{\text{Im}\Pi_{mol}(s)}{s - q^2 - i\epsilon} + ... \tag{3}$$

where $\text{Im}\Pi_{mol}(s)$ are the spectral functions.

• *Laplace sum rule (LSR)*
The Laplace SVZ sum rules are improvement of the previous dispersion relation which becomes after the (inverse) Laplace transform:

$$\mathcal{L}(\tau) = \int_{4m_Q^2}^{\infty} dt\, e^{-t\tau}\frac{1}{\pi}\text{Im}\Pi^{OPE}_{mol}(t)\,. \tag{4}$$

Parametrizing the spectral function by one resonance plus a QCD continuum, the lowest resonance mass M_H and coupling f_H normalized as:

$$\langle 0|J^\mu|H\rangle = f_H M_H^4 \epsilon^\mu\,, \tag{5}$$

can be extracted from the previous Laplace sum rules (LSR) as [4, 5]:

$$M_H^2 = \frac{\int_{4m_Q^2}^{t_c} dt\, t\, e^{-t\tau}\frac{1}{\pi}\text{Im}\Pi^{OPE}_{mol}(t)}{\int_{4m_Q^2}^{t_c} dt\, e^{-t\tau}\frac{1}{\pi}\text{Im}\Pi^{OPE}_{mol}(t)} \tag{6}$$

and

$$f_H^2 = \frac{\int_{4m_Q^2}^{t_c} dt\, e^{-t\tau}\frac{1}{\pi}\text{Im}\Pi^{OPE}_{mol}(t)}{e^{-\tau M_H^2} M_H^8}\,, \tag{7}$$

*Talk given at QCD 14 (29 june - 3 july, Montpellier - France).

**Speaker.
Email addresses: fanfenos@yahoo.fr (F. Fanomezana), snarison@yahoo.fr (S. Narison), achris_01@yahoo.fr (Andry Rabemananjara)

where M_Q is the heavy quark mass, τ the sum rule parameter and t_c the continuum threshold.

• *The QCD two-point function at N2LO*

For evaluating the perturbative part at NLO and N2LO, we assume a factorization of the bilinear currents and do a convolution of the corresponding scalar $\text{Im}\Pi^{(0)}(s_1)$ and vector $\text{Im}\Pi^{(1)}(s_2)$ correlators [6, 7]:

$$\frac{1}{\pi}\text{Im}\Pi^{(1)}_{mol}(t) = \theta(t-4M_Q^2)\left(\frac{1}{4\pi}\right)^2 t^2 \int_{M_Q^2}^{(\sqrt{t}-M_Q)^2} dt_1 \times$$
$$\int_{M_Q^2}^{(\sqrt{t}-\sqrt{t_1})^2} dt_2\, \lambda^{3/2}\frac{1}{\pi}\text{Im}\Pi^{(1)}(t_1)\frac{1}{\pi}\text{Im}\Pi^{(0)}(t_2) \qquad (8)$$

with the phase space factor:

$$\lambda = \left(1-\frac{(\sqrt{t}-\sqrt{t_1})^2}{t}\right)\left(1-\frac{(\sqrt{t_1}+\sqrt{t_2})^2}{t}\right). \qquad (9)$$

Their QCD expressions are known in the literature [8–11]. We shall use the relation between the on-shell M_Q and the running mass $\bar{m}_Q(\nu)$ to transform the spectral function into the $\overline{MS}$-scheme [12, 13]:

$$M_Q = \overline{m}_Q(\nu)\Bigg[1+\frac{4}{3}a_s + (16.2163-1.0414n_l)a_s^2$$
$$+\ln\left(\frac{\nu}{M_Q}\right)^2\left(a_s+(8.8472-0.3611n_l)a_s^2\right)$$
$$+\ln^2\left(\frac{\nu}{M_Q}\right)^2(1.7917-0.0833n_l)\,a_s^2\Bigg], \qquad (10)$$

where $n_l = n_f - 1$ is the number of light flavours and $a_s(\nu) = \alpha_s(\nu)/\pi$ at the scale ν. The QCD expressions of the non-perturbative part of the 1^{--} molecule spectral functions used here have been computed up to dimension $d = 8$ in e.g. Ref. [14].

• *QCD parameters*

The PT QCD parameters which appear in this analysis are α_s, the charm and bottom quark masses $m_{c,b}$ (the light quark masses have been neglected). The non-perturbative condensates up to dimension 8 considered here are the quark condensate $\langle\bar{q}q\rangle$, the two-gluon condensate $\langle g^2G^2\rangle$, the mixed condensate $\langle g\bar{q}Gq\rangle$, the four-quark condensate $\rho\langle\bar{q}q\rangle^2$, the three-gluon condensate $\langle g^3G^3\rangle$, and the two-quark multiply two-gluon condensate $\rho\langle\bar{q}q\rangle\langle g^2G^2\rangle$ where ρ indicates the deviation from the four-quark vacuum saturation. Their values are given in Table 1. For the condensates, we shall use:

$$\langle\bar{q}q\rangle(\nu) = -\hat{\mu}_q^3\left(\text{Log}\frac{\nu}{\Lambda}\right)^{-2/\beta_1} \qquad (11)$$

$$\langle g\bar{q}Gq\rangle(\nu) = -M_0^2\hat{\mu}_q^3\left(\text{Log}\frac{\nu}{\Lambda}\right)^{-1/3\beta_1} \qquad (12)$$

where $\beta_1 = -(1/2)(11-2n_f/3)$ is the first coefficient of the β function, $\hat{\mu}_q$ the renormalization group invariant condensate and Λ is the QCD scale.

Parameters	Values.
$\alpha_s(M_\tau)$	0.325(8)
$\Lambda(n_f=4)$	(324 ± 15) MeV
$\Lambda(n_f=5)$	(194 ± 10) MeV
$\bar{m}_c(m_c)$	(1261 ± 24) MeV
$\bar{m}_b(m_b)$	(4177 ± 22) MeV
$\hat{\mu}_q$	(263 ± 7) MeV
M_0^2	(0.8 ± 0.2) GeV^2
$\langle\alpha_sG^2\rangle$	$(7\pm2)\times10^{-2}$ GeV^4
$\langle g^3G^3\rangle$	(8.2 ± 2.0) GeV^2 × $\langle\alpha_sG^2\rangle$
$\rho = \langle\bar{q}q\bar{q}q\rangle/\langle\bar{q}q\rangle^2$	(2 ± 1)

Table 1: QCD input parameters (see e.g. [5, 15] and references therein).



Figure 1: a) τ-behaviour of the $M_{D^*D_0^*}$ for different values of t_c and for $\nu = 4m_c$; the same as in a) but for the running coupling $f_{D^*D_0^*}$.

3. $\bar{D}^*D_0^*(1^{--})$ molecule mass and coupling

• *τ and t_c stabilities*

We show in Fig. 1a the τ−behaviour of the $M_{\bar{D}^*D_0^*}$ at N2LO of PT series for different values of t_c at given subtraction $\nu = 4m_c$ and in Fig. 1b the one of the coupling $f_{\bar{D}^*D_0^*}$, where one should note that the bilinear scalar heavy-light current acquires an anomalous dimension such that the decay constant runs as:

$$f_{\bar{D}^*D_0^*}(\nu) = \hat{f}_{\bar{D}^*D_0^*}\left(\text{Log}\frac{\nu}{\Lambda}\right)^{2/-\beta_1}, \qquad (13)$$

where $\hat{f}_{\bar{D}^*D_0^*}$ is a scale invariant coupling. One can see in these figures that the τ stability is obtained from $t_c = 32 \sim 36$ GeV^2 while t_c stability is reached from $t_c = 49$ GeV^2. We consider as optimal and conservative result the one obtained inside this region of t_c.



Figure 2: a) $M_{\bar{D}^* D_0^*}$ mass versus τ for two extremal values of t_c = 36 and 49 GeV^2 and for different truncation of the PT series; b) the as in a) but for but for the running coupling $f_{\bar{D}^* D_0}$.

• *Convergence of the PT series*

We show in the Fig. 2, the convergence of the expansion in PT. We can observe in Fig. 2a that, from the LO to the NLO, the mass increases by +6.72% and from LO $\oplus$ NLO to N2LO, by +4.42%. These corrections indicate that the LO results obtained in [3, 14], by using the value of the running m_c mass, underestimates the molecule mass by about 11.14%.

For the coupling, the α_s corrections are large. It increases about 50% from LO to NLO and about 35% from LO $\oplus$ NLO to N2LO, then to a total of 85%. Nevertheless, one can see that the PT series still converges though slowly.

• *ν-stability*

We study in Fig. 3 the behaviour of the mass and of the invariant coupling $\hat{f}_{\bar{D}^* D_0^*}$ in term of the scale ν. One notices a good stability for ν ranging from 2.5 to 6 GeV where a minimum for the mass and an inflexion for the coupling occur for $\nu \approx 3m_c = 3.8$ GeV which the same value as the one in the 1^{++} channel. We obtain at N2LO:

$$M_{\bar{D}^* D_0^*} = 5244(228)\ \text{MeV}\ , \tag{14}$$

which is comparable with the LO result $M_{\bar{D}^* D_0^*} \simeq 5268(24)$ MeV obtained by combining LSR and FESR [3]. For the coupling, we find:

$$\begin{aligned} \hat{f}_{\bar{D}^* D_0^*} &= (13.7 \pm 0.9) \times 10^{-2}\text{MeV} \implies \\ f_{\bar{D}^* D_0^*}(\nu) &= (21.3 \pm 1.4) \times 10^{-2}\text{MeV}\ , \end{aligned} \tag{15}$$

which we compare with the LO result $f_{\bar{D}^* D_0^*} \approx 0.08$ MeV obtained in [3] for a 1^{--} four-quark state where the difference between the two values comes mainly from the large radiative corrections discussed before.



Figure 3: a) ν-behaviour of the $M_{\bar{D}^* D_0^*}$; b) the same as in a) but for $M_{\bar{D}^* D_0^*}$.

4. $\bar{B}^* B_0(1^{--})$ mass and coupling

• *τ and t_c stabilities*

We do the same analysis for the case of the beauty channel. The $\tau-$behavior of mass and running coupling is shown in the Fig. 4 for given value of ν and for different values of t_c. One can see that the stability in τ is between t_c = 135 GeV^2 and t_c = 160 GeV^2 where t_c stability starts. We consider as optimal and conservative value of the mass and coupling the one obtained between these two extremal cases.

• *Convergence of the PT series*

We study in Fig. 5 the convergence of the PT series. From LO to the NLO, the mass increases by 4.9% and from LO $\oplus$ NLO to N2LO it decreases by -2.98% which shows small corrections and a good convergence of the PT series. For the coupling, the α_s correction increases the value by about 25% while from LO $\oplus$ NLO to N2LO, its decreases by about -15%.

• *ν-stability*

We study in Fig. 6 the behaviour of the mass and of the invariant coupling $\hat{f}_{\bar{B}^* B_0^*}$ in term of the scale ν. Our final result for the mass comes from $\nu \approx 3.5$ GeV where stabilities are obtained both for the mass and for the couling. We obtain:

$$M_{\bar{B}^* B_0^*} = 11920(159)\ \text{MeV}\ . \tag{16}$$

and:

$$\begin{aligned} \hat{f}_{\bar{B}^* B_0^*} &= 2.0(0.59) \times 10^{-2}\ \text{MeV} \implies \\ f_{\bar{B}^* B_0^*}(\nu) &= 3.5(1.02) \times 10^{-2}\ \text{MeV}\ . \end{aligned} \tag{17}$$



Figure 4: a) τ-behaviour of the $M_{\bar{B}^* B_0^*}$ for different t_c and for $\nu = m_b$; the same as in a) but for the running coupling $f_{\bar{B}^* B_0^*}$.



Figure 5: a) $M_{\bar{B}^* B_0^*}$ mass versus τ for two extremal values of t_c = 36 and 49 GeV^2 and for different truncation of the PT series; b) the as in a) but for but for the running coupling $f_{\bar{B}^* B_0}$.

5. Conclusions

We have studied the $\bar{D}^* D_0^*$ and $\bar{B}^* B_0^*$ (1^{--}) molecules states using QCD spectral sum rules to N2LO of PT series and including non-perturbative condensates up to dimension 8. We consider the present result as an improvement of the previous ones [3, 14].

The mass of $\bar{D}^* D_0^*$ in Eq. (14) at N2LO agrees within the error with the LO result [3, 14] due to the small radiative corrections in the ratio of moments, as expected from a general feature of this approach. The result shows that the 1^{--} experimental candidates Y(4260), Y(4360), Y(4660) are too light to be pure $\bar{D}^* D_0^*$ molecule states.

The coupling $f_{\bar{D}^* D_0^*}$ obtained in Eq. (15) is strongly affected by radiative corrections and using its LO value for estimating some hadronic widths may lead to very inaccurate predictions.

For the beauty channel, our predictions in Eq. (16) can be checked in different B factories. Here, the radiative corrections are smaller for the coupling.



Figure 6: a) ν-behaviour of the $M_{\bar{B}^* B_0^*}$; b) the same as in a) but for $f_{\bar{B}^* B_0^*}$.

Acknowledgements

F.F. and A.R. would like to thank the CNRS for supporting the travel and living expenses and the LUPM-Montpellier for hospitality. We also thank R.M. Albuquerque for many helpful discussions.

References

[1] B. Aubert et al. (BaBar Collaboration), *Phys. Rev. Lett.* **95** (2005) 142001; Q. He et al.(CLEO Collaboration), *Phys. Rev.* **D74** (2006) 091104(R); C. Z. Yuan et al. (Belle Collaboration), *Phys. Rev. Lett.* **99** (2007) 182004; T. E. Coan et al. (CLEO Collaboration), *Phys. Rev. Lett.* **96** (2006) 162003; J. P. Lees et al. (BaBar Collaboration), *Phys. Rev.* **D86** (2012) 051102(R).

[2] For reviews, see e.g.: F. S. Navarra, M. Nielsen, S. H. Lee, *Phys. Rep.* **497** (2010) 4; S. L. Zhu, *Int. J. Mod. Phys.* **E17** (2008); E.Swanson, *Phys. Rep.* **429** (2006) 243; N. Brambilla et al., *Eur. Phys. J* **C71** (2011) 1534.

[3] R.M. Albuquerque, F. Fanomezana, S. Narison, A. Rabemananjara, *Phys. Lett.* **B 715** (2012) 129.

[4] M.A. Shifman, A.I. and Vainshtein and V.I. Zakharov, Nucl. Phys. **B147** (1979) 385 .

[5] For a review and references to original works, see e.g., S. Narison, *QCD as a theory of hadrons, Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol.* **17** (2002) 1-778 [hep-h/0205006].

[6] A. Pich and E. de Rafael, *Phys. Lett.***B158** (1985) 477.

[7] S. Narison and A. Pivovarov, *Phys. Lett.* **B327** (1994) 341.

[8] S. Narison, *Phys. Lett.* **B718** (2013) 1321.

[9] S. Narison, arXiv:1404.6642 [hep-ph].

[10] S.C. Generalis, Ph.D. thesis, Open Univ. report, OUT-4102-13 (1982), unpublished.

[11] K.G. Chetyrkin and M. Steinhauser, *Phys. Lett.* **B 502** (2001) 104; hep-ph/0108017.

[12] S. Narison, *Phys. Lett.* **B197** (1987) 405; *Phys. Lett.***B 216** (1989) 191.

[13] K.G. Chetyrkin and M. Steinhauser, *Nucl. Phys.* **B 573** (2000) 617; K. Melnikov and T. van Ritbergen, hep-ph/9912391.

[14] R. M. Albuquerque and M. Nielsen, *Nucl. Phys.* **A815** (2009) 53; Erratum-ibid. **A857** (2011) 48 and private notes from R. Albuquerque.

[15] S. Narison, arXiv:1409.8148 (talk given at this conference).